

DFTT 68/97
 IASSNS-AST 97/66
 UWThPh-1997-44
 hep-ph/9711416

Neutrino masses and mixing from neutrino oscillation experiments

S.M. Bilenky

*Joint Institute for Nuclear Research, Dubna, Russia, and
 Institute for Advanced Study, Princeton, N.J. 08540*

C. Giunti

*INFN, Sezione di Torino, and Dipartimento di Fisica Teorica, Università di Torino,
 Via P. Giuria 1, I-10125 Torino, Italy*

W. Grimus

*Institute for Theoretical Physics, University of Vienna,
 Boltzmanngasse 5, A-1090 Vienna, Austria*

Abstract

We discuss which information on neutrino masses and mixing can be obtained from the results of all neutrino oscillation experiments in the cases of three and four massive neutrinos. We show that in the three-neutrino case the neutrino oscillation data are not compatible with a hierarchy of couplings. In the case of four neutrinos, a hierarchy of masses is disfavored by the data and only two schemes with two pairs of neutrinos with close masses separated by a gap of the order of 1 eV can accommodate the results of all experiments.

Talk presented by C. Giunti at the *International Europhysics Conference on High Energy Physics*, 19–26 August 1997, Jerusalem, Israel.

Typeset using REVTeX

The search for neutrino oscillations is one of the most active branches of today's high-energy physics. From the LEP measurements of the invisible width of the Z -boson we know that there are three light active flavor neutrinos: ν_e , ν_μ and ν_τ . In general, flavor neutrinos are not mass eigenstates and the left-handed flavor neutrino fields $\nu_{\alpha L}$ are superpositions of the left-handed components ν_{kL} of the fields of neutrinos with a definite mass ($k = 1, 2, 3, \dots, n$): $\nu_{\alpha L} = \sum_{k=1}^n U_{\alpha k} \nu_{kL}$, where U is a unitary mixing matrix. The number n of massive neutrinos can be three or more, without any experimental upper limit. If $n > 3$, there are $n - 3$ sterile flavor neutrino fields, i.e., fields of neutrinos which do not take part in standard weak interactions; in this case $\nu_\alpha = \nu_e, \nu_\mu, \nu_\tau, \nu_{s1}, \nu_{s2}, \dots, \nu_{sn-3}$. Neutrino oscillations is a direct consequence of neutrino mixing; the probability of $\nu_\alpha \rightarrow \nu_\beta$ transitions is given by (see [1])

$$P_{\nu_\alpha \rightarrow \nu_\beta} = \left| \sum_{k=1}^n U_{\beta k} U_{\alpha k}^* \exp \left(-i \frac{\Delta m_{k1}^2 L}{2E} \right) \right|^2, \quad (1)$$

where L is the distance between the neutrino source and detector, E is the neutrino energy and $\Delta m_{kj}^2 \equiv m_k^2 - m_j^2$.

In this report we discuss which information on the neutrino mass spectrum and mixing parameters can be obtained from the results of neutrino oscillation experiments. Many short-baseline (SBL) neutrino oscillation experiments with reactor and accelerator neutrinos did not find any evidence of neutrino oscillations. Their results can be used in order to constrain the allowed values of the neutrino masses and of the elements of the mixing matrix. In our analysis we use the most stringent exclusion plots obtained in the $\bar{\nu}_e \rightarrow \bar{\nu}_e$ channel by the Bugey experiment [2], in the $\nu_\mu \rightarrow \nu_\mu$ channel by the CDHS and CCFR experiments [3], and in the $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ channel by the BNL E734, BNL E776 and CCFR [4] experiments.

There are three experimental indications in favor of neutrino oscillations that come from the anomalies observed by the solar neutrino experiments [5], the atmospheric neutrino experiments [6] and the LSND experiment [7]. The solar neutrino deficit can be explained with oscillations of solar ν_e 's into other states and indicates a mass-squared difference of the order of 10^{-5} eV^2 in the case of resonant MSW transitions or 10^{-10} eV^2 in the case of vacuum oscillations. The atmospheric neutrino anomaly can be explained by $\bar{\nu}_\mu \rightarrow \bar{\nu}_x$ oscillations ($x \neq \mu$) with a mass-squared difference of the order of 10^{-2} eV^2 . Finally, the LSND experiment found indications in favor of $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations with a mass-squared difference of the order of 1 eV^2 .

Hence, three different scales of mass-squared difference are needed in order to explain the three indications in favor of neutrino oscillations. This means that the number of massive neutrinos must be bigger than three. In the following we consider the simplest possibility, i.e. the existence of four massive neutrinos ($n = 4$). In this case, besides the three light flavor neutrinos ν_e, ν_μ, ν_τ , there is a light sterile neutrino ν_s .

However, before considering the case of four neutrinos, we discuss the minimal possibility of the existence of only three massive neutrinos ($n = 3$). In this case one of the experimental anomalies mentioned above cannot be explained with neutrino oscillations (we choose to disregard the atmospheric neutrino anomaly).

In both cases of three and four massive neutrinos the oscillations in the LSND experiment imply that the largest mass-squared difference $\Delta m_{n1}^2 \equiv m_n^2 - m_1^2$ is relevant for

SBL oscillations, whereas the other mass-squared differences are much smaller. Hence, the mass spectrum must be composed of two groups of massive neutrinos with close masses ($\nu_1, \dots, \nu_{r-1}$ and $\nu_r, \dots, \nu_n$) separated by a mass difference in the eV range ($m_1 < \dots < m_{r-1} \ll m_r < \dots < m_n$) and in SBL experiments we have $\frac{\Delta m_{n1}^2 L}{2E} \gtrsim 1$, $\frac{\Delta m_{k1}^2 L}{2E} \ll 1$ for $k < r$, $\frac{\Delta m_{nk}^2 L}{2E} \ll 1$ for $k \geq r$. The formula (1) written as

$$P_{\nu_\alpha \rightarrow \nu_\beta} = \left| \sum_{k=1}^{r-1} U_{\beta k} U_{\alpha k}^* e^{-i \frac{\Delta m_{k1}^2 L}{2E}} + e^{-i \frac{\Delta m_{n1}^2 L}{2E}} \sum_{k=r}^n U_{\beta k} U_{\alpha k}^* e^{i \frac{\Delta m_{nk}^2 L}{2E}} \right|^2 \quad (2)$$

leads to the following expression for the transition ($\beta \neq \alpha$) and survival ($\beta = \alpha$) probabilities of neutrinos and anti-neutrinos in SBL experiments:

$$P_{\nu_\alpha \rightarrow \nu_\beta}^{(\text{SBL})} = A_{\alpha;\beta} \sin^2 \frac{\Delta m^2 L}{4E}, \quad P_{\nu_\alpha \rightarrow \nu_\alpha}^{(\text{SBL})} = 1 - B_{\alpha;\alpha} \sin^2 \frac{\Delta m^2 L}{4E}, \quad (3)$$

with $\Delta m^2 \equiv \Delta m_{n1}^2$ and the oscillation amplitudes

$$A_{\alpha;\beta} = 4 \left| \sum_{k=r}^n U_{\beta k} U_{\alpha k}^* \right|^2, \quad B_{\alpha;\alpha} = 4 \left(\sum_{k=r}^n |U_{\alpha k}|^2 \right) \left(1 - \sum_{k=r}^n |U_{\alpha k}|^2 \right). \quad (4)$$

The formulas (3) have the same form of the standard expressions for the oscillation probabilities in the case of two neutrinos (see [1]) with which the data of all the SBL experiments have been analyzed by the experimental groups. Hence, the results of these analyses can be used in order to constrain the possible values of the oscillation amplitudes $A_{\alpha;\beta}$ and $B_{\alpha;\alpha}$.

First, we consider the scheme 3H of Tab.I, with three neutrinos and a mass hierarchy. This scheme (as all the schemes with three neutrinos) provides only two independent mass-squared differences, Δm_{21}^2 and Δm_{31}^2 , which we choose to be relevant for the solution of the solar neutrino problem and for neutrino oscillations in the LSND experiment, respectively.

Let us emphasize that the mass spectrum 3H with three neutrinos and a mass hierarchy is the simplest and most natural one, being analogous to the mass spectra of charged leptons, up and down quarks. Moreover, a scheme with three neutrinos and a mass hierarchy is predicted by the see-saw mechanism for the generation of neutrino masses, which can explain the smallness of the neutrino masses with respect to the masses of the corresponding charged leptons.

In the case of scheme 3H we have $n = r = 3$ and Eq.(4) implies that

$$A_{\alpha;\beta} = 4 |U_{\alpha 3}|^2 |U_{\beta 3}|^2, \quad B_{\alpha;\alpha} = 4 |U_{\alpha 3}|^2 (1 - |U_{\alpha 3}|^2). \quad (5)$$

Hence, neutrino oscillations in SBL experiments depend on three parameters: $\Delta m^2 \equiv \Delta m_{31}^2$, $|U_{e3}|^2$ and $|U_{\mu 3}|^2$ (the unitarity of U implies that $|U_{\tau 3}|^2 = 1 - |U_{e3}|^2 - |U_{\mu 3}|^2$). From the exclusion plots obtained in reactor $\bar{\nu}_e$ and accelerator ν_μ disappearance experiments it follows that at any fixed value of Δm^2 , the oscillation amplitudes $B_{e;e}$ and $B_{\mu;\mu}$ are bounded by the upper values $B_{e;e}^0$ and $B_{\mu;\mu}^0$, respectively, which are small quantities for $0.3 \lesssim \Delta m^2 \lesssim 10^3 \text{ eV}^2$. From Eq.(5) one can see that small upper bounds for $B_{e;e}$ and $B_{\mu;\mu}$ imply that the parameters $|U_{e3}|^2$ and $|U_{\mu 3}|^2$ can be either small or large (i.e., close to one):

$$|U_{\alpha 3}|^2 \leq a_\alpha^0 \quad \text{or} \quad |U_{\alpha 3}|^2 \geq 1 - a_\alpha^0, \quad \text{with} \quad a_\alpha^0 = \frac{1}{2} \left(1 - \sqrt{1 - B_{\alpha;\alpha}^0} \right), \quad (6)$$

for $\alpha = e, \mu$. Both a_e^0 and a_μ^0 are small ($a_e^0 \lesssim 4 \times 10^{-2}$ and $a_\mu^0 \lesssim 2 \times 10^{-1}$) for any value of Δm^2 in the range $0.3 \lesssim \Delta m^2 \lesssim 10^3 \text{ eV}^2$ (see Fig.1 of Ref. [8]).

Since large values of both $|U_{e3}|^2$ and $|U_{\mu 3}|^2$ are excluded by the unitarity of the mixing matrix ($|U_{e3}|^2 + |U_{\mu 3}|^2 \leq 1$), at any fixed value of Δm^2 there are three regions in the $|U_{e3}|^2$ – $|U_{\mu 3}|^2$ plane which are allowed by the exclusion plots of SBL disappearance experiments: Region I, with $|U_{e3}|^2 \leq a_e^0$ and $|U_{\mu 3}|^2 \leq a_\mu^0$; Region II, with $|U_{e3}|^2 \leq a_e^0$ and $|U_{\mu 3}|^2 \geq 1 - a_\mu^0$; Region III, with $|U_{e3}|^2 \geq 1 - a_e^0$ and $|U_{\mu 3}|^2 \leq a_\mu^0$.

In region III $|U_{e3}|^2$ is large and ν_e has a small mixing with ν_1 and ν_2 , which is insufficient for the explanation of the solar neutrino problem. Indeed, the survival probability of solar ν_e 's is bounded by $P_{\nu_e \rightarrow \nu_e}^{\text{sun}} \geq |U_{e3}|^4$ (see [8]). If $|U_{e3}|^2 \geq 1 - a_e^0$, we have $P_{\nu_e \rightarrow \nu_e}^{\text{sun}} \gtrsim 0.92$ at all neutrino energies, which is a bound that is not compatible with the solar neutrino data. Hence, region III is excluded by solar neutrinos.

In region I $A_{\mu;e} \leq 4 a_e^0 a_\mu^0$, which means that the probability of $\bar{\nu}_\mu \leftrightarrow \bar{\nu}_e$ transitions in SBL experiments is strongly suppressed. The corresponding upper bound obtained from the 90% CL exclusion plots of the Bugey [2] $\bar{\nu}_e$ disappearance experiment and of the CDHS and CCFR [3] ν_μ disappearance experiments is represented in Fig.1 by the curve passing through the circles. The shadowed regions in Fig.1 are allowed at 90% CL by the results of the LSND experiment. Also shown are the 90% CL exclusion curves found in the BNL E734, BNL E776 and CCFR [4] $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ appearance experiments and in the Bugey experiment. One can see from Fig.1 that in region I, the bounds obtained from the results of $\bar{\nu}_e \rightarrow \bar{\nu}_e$, $\nu_\mu \rightarrow \nu_\mu$ and $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ experiments are not compatible with the allowed regions of the LSND experiment [8]. Therefore, region I is disfavored by the results of SBL experiments. This is an important indication, because region I is the only one in which it is possible to have a hierarchy of the elements of the neutrino mixing matrix analogous to the one of the quark mixing matrix.

Having excluded the regions I and III of the scheme 3H, we are left only with region II, where ν_μ has a large mixing with ν_3 , i.e., ν_μ (not ν_τ) is the “heaviest” neutrino.

Let us now consider the possible schemes with four neutrinos, which provide three independent mass-squared differences and allow to accommodate in a natural way all the three experimental indications in favor of neutrino oscillations. We consider first the scheme 4H of Tab.I with four neutrinos and a mass hierarchy. The three independent mass-squared differences, Δm_{21}^2 , Δm_{31}^2 and Δm_{41}^2 , are taken to be relevant for the oscillations of solar, atmospheric and LSND neutrinos, respectively. In the case of scheme 4H we have $n = r = 4$ and Eq.(4) implies that the oscillation amplitudes are given by

$$A_{\alpha;\beta} = 4 |U_{\alpha 4}|^2 |U_{\beta 4}|^2, \quad B_{\alpha;\alpha} = 4 |U_{\alpha 4}|^2 (1 - |U_{\alpha 4}|^2). \quad (7)$$

In this case, neutrino oscillations in SBL experiments depend on four parameters: $\Delta m^2 \equiv \Delta m_{31}^2$, $|U_{e3}|^2$, $|U_{\mu 3}|^2$ and $|U_{\tau 3}|^2$. From the similarity of the amplitudes (7) with the corresponding ones given in Eq.(5), it is clear that replacing $|U_{\alpha 3}|^2$ with $|U_{\alpha 4}|^2$ we can apply to the scheme 4H the same analysis presented for the scheme 3H. Hence, also the regions III and I of the scheme 4H are excluded, respectively, by the solar neutrino problem and by the

results of SBL experiments. Furthermore, the purpose of considering the scheme 4H is to have the possibility to explain the atmospheric neutrino anomaly, but this is not possible if the neutrino mixing parameters lie in region II. Indeed, in region II $|U_{\mu 4}|^2$ is large and the muon neutrino has a small mixing with the light neutrinos ν_1 , ν_2 and ν_3 , which is insufficient for the explanation of the atmospheric neutrino anomaly [9].

Hence, the scheme 4H is disfavored by the results of neutrino oscillation experiments. For the same reasons, all possible schemes with four neutrinos and a mass spectrum in which three masses are clustered and one mass is separated from the others by a gap of about 1 eV (needed for the explanation of the LSND data) are disfavored by the results of neutrino oscillation experiments. Therefore, there are only two possible schemes with four neutrinos which are compatible with the results of all neutrino oscillation experiments: the schemes 4A and 4B of Tab.I. In these two schemes the four neutrino masses are divided in two pairs of close masses separated by a gap of about 1 eV. In scheme A, Δm_{21}^2 is relevant for the explanation of the atmospheric neutrino anomaly and Δm_{43}^2 is relevant for the suppression of solar ν_e 's. In scheme B, the roles of Δm_{21}^2 and Δm_{43}^2 are reversed.

From Eq.(4) and using the unitarity of the mixing matrix, the oscillation amplitudes $B_{\alpha;\alpha}$ in the schemes 4A and 4B ($n = 4$, $r = 3$) are given by $B_{\alpha;\alpha} = 4c_\alpha(1 - c_\alpha)$, with $c_\alpha \equiv \sum_{k=1,2} |U_{\alpha k}|^2$ in the scheme 4A and $c_\alpha \equiv \sum_{k=3,4} |U_{\alpha k}|^2$ in the scheme 4B. This expression for $B_{\alpha;\alpha}$ has the same form as the one in Eq.(7), with $|U_{\alpha 4}|^2$ replaced by c_α . Therefore, we can apply the same analysis presented for the scheme 4H and we obtain four allowed regions in the c_e - c_μ plane (now the region with large c_e and c_μ is not excluded by the unitarity of the mixing matrix, which gives the constraint $c_e + c_\mu \leq 2$): Region I, with $c_e \leq a_e^0$ and $c_\mu \leq a_\mu^0$; Region II, with $c_e \leq a_e^0$ and $c_\mu \geq 1 - a_\mu^0$; Region III, with $c_e \geq 1 - a_e^0$ and $c_\mu \leq a_\mu^0$; Region IV, with $c_e \geq 1 - a_e^0$ and $c_\mu \geq 1 - a_\mu^0$. Following the same reasoning as in the case of scheme 4H, one can see that the regions III and IV are excluded by the solar neutrino data and the regions I and III are excluded by the results of the atmospheric neutrino experiments [9]. Hence, only region II is allowed by the results of all experiments.

If the neutrino mixing parameters lie in region II, in the scheme 4A (4B) the electron (muon) neutrino is “heavy”, because it has a large mixing with ν_3 and ν_4 , and the muon (electron) neutrino is “light”. Thus, the schemes 4A and 4B give different predictions for the effective Majorana mass $\langle m \rangle = \sum_k U_{ek}^2 m_k$ in neutrinoless double-beta decay experiments: since $m_3 \simeq m_4 \gg m_1 \simeq m_2$, we have $|\langle m \rangle| \leq (1 - c_e)m_4 \simeq m_4$ in the scheme 4A and $|\langle m \rangle| \leq c_e m_4 \leq a_e^0 m_4 \ll m_4$ in the scheme 4B. Hence, if the scheme 4A is realized in nature, the experiments on the search for neutrinoless double-beta decay can reveal the effects of the heavy neutrino masses $m_3 \simeq m_4$. Furthermore, the smallness of c_e in both schemes 4A and 4B implies that the electron neutrino has a small mixing with the neutrinos whose mass-squared difference is responsible for the oscillations of atmospheric neutrinos (i.e., ν_1 , ν_2 in scheme 4A and ν_3 , ν_4 in scheme 4B). Hence, the transition probability of electron neutrinos and antineutrinos into other states in atmospheric and long-baseline experiments is suppressed [10]¹.

¹After we finished this paper the results of the first long-baseline reactor experiment CHOOZ appeared (M. Apollonio *et al.*, preprint hep-ex/9711002). No indications in favor of $\bar{\nu}_e \rightarrow \bar{\nu}_e$

REFERENCES

- [1] S.M. Bilenky and B. Pontecorvo, Phys. Rep. **41**, 225 (1978).
- [2] B. Achkar *et al.*, Nucl. Phys. B **434**, 503 (1995).
- [3] F. Dydak *et al.*, Phys. Lett. B **134**, 281 (1984); I.E. Stockdale *et al.*, Phys. Rev. Lett. **52**, 1384 (1984).
- [4] L.A.Ahrens *et al.*, Phys. Rev. D **36**, 702 (1987); L.Borodovsky *et al.*, Phys. Rev. Lett. **68**, 274 (1992); A.Romosán *et al.*, *ibid* **78**, 2912 (1997).
- [5] B.T. Cleveland *et al.*, Nucl. Phys. B (P.S.) **38**, 47 (1995); K.S. Hirata *et al.*, Phys. Rev. D **44**, 2241 (1991); GALLEX Coll., Phys. Lett. B **388**, 384 (1996); SAGE Coll., Phys. Rev. Lett. **77**, 4708 (1996).
- [6] Y. Fukuda *et al.*, Phys. Lett. B **335**, 237 (1994); R. Becker-Szendy *et al.*, Nucl. Phys. B (P.S.) **38**, 331 (1995); W.W.M. Allison *et al.*, Phys. Lett. B **391**, 491 (1997).
- [7] C. Athanassopoulos *et al.*, Phys. Rev. Lett. **77**, 3082 (1996).
- [8] S.M. Bilenky, A. Bottino, C. Giunti and C.W. Kim, Phys. Rev. D **54**, 1881 (1996).
- [9] S.M. Bilenky, C. Giunti and W. Grimus, preprint hep-ph/9607372.
- [10] S.M. Bilenky, C. Giunti and W. Grimus, preprint hep-ph/9710209.

transitions were found in this experiment. The upper bound for the transition probability of electron antineutrinos into other states found in the CHOOZ experiment is in agreement with the limit obtained in Ref. [10].

$$3\text{H: } \overbrace{m_1 \ll m_2 \ll m_3}^{\text{sun}} \\ \text{LSND}$$

$$4\text{H: } \overbrace{\overbrace{m_1 \ll m_2 \ll m_3 \ll m_4}^{\text{sun}}}^{\text{atm}} \\ \text{LSND}$$

$$4\text{A: } \overbrace{m_1 < m_2 \ll m_3 < m_4}^{\text{atm}} \quad \overbrace{\quad}^{\text{sun}} \\ \text{LSND}$$

$$4\text{B: } \overbrace{m_1 < m_2 \ll m_3 < m_4}^{\text{sun}} \quad \overbrace{\quad}^{\text{atm}} \\ \text{LSND}$$

Table I




Figure 1